

Cyclic codes over $\mathbb{F}_{2^m}[u]/\langle u^k \rangle$ of oddly even length

Yonglin Cao¹ · Yuan Cao² · Fang-Wei Fu³

Received: date / Accepted: date

Abstract Let $\mathbb{F}_{2^m}$ be a finite field of characteristic 2 and $R = \mathbb{F}_{2^m}[u]/\langle u^k \rangle = \mathbb{F}_{2^m} + u\mathbb{F}_{2^m} + \dots + u^{k-1}\mathbb{F}_{2^m}$ ($u^k = 0$) where $k \in \mathbb{Z}^+$ satisfies $k \geq 2$. For any odd positive integer n , it is known that cyclic codes over R of length $2n$ are identified with ideals of the ring $R[x]/\langle x^{2n} - 1 \rangle$. In this paper, an explicit representation for each cyclic code over R of length $2n$ is provided and a formula to count the number of codewords in each code is given. Then a formula to calculate the number of cyclic codes over R of length $2n$ is obtained. Moreover, the dual code of each cyclic code and self-dual cyclic codes over R of length $2n$ are investigated.

Keywords Cyclic code · Finite chain ring · Non-principal ideal ring · Dual code · Self-dual code

Mathematics Subject Classification (2000) 94B05 · 94B15 · 11T71

1 Introduction and preliminaries

Throughout this paper, let $\mathbb{F}_{p^m}$ be a finite field of characteristic p and cardinality p^m , and denote $R = \mathbb{F}_{p^m}[u]/\langle u^k \rangle = \mathbb{F}_{p^m} + u\mathbb{F}_{p^m} + \dots + u^{k-1}\mathbb{F}_{p^m}$ ($u^k = 0$), where p is a prime and $k \in \mathbb{Z}^+$ satisfying $k \geq 2$. It is well-known that R is a finite chain ring. For any positive integer N , let $R[x]/\langle x^N - 1 \rangle$ be the residue class ring of the polynomial ring $R[x]$ modulo its ideal $\langle x^N - 1 \rangle$ generated by

Yonglin Cao
 E-mail: ylcao@sdut.edu.cn
 Yuan Cao
 E-mail: yuan-cao@hnu.edu.cn
 Fang-Wei Fu
 E-mail: fwfu@nankai.edu.cn

¹ School of Sciences, Shandong University of Technology, Zibo, Shandong 255091, China

² College of Mathematics and Econometrics, Hunan University, Changsha 410082, China

³ Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, China

$x^N - 1$. Then every element of $R[x]/\langle x^N - 1 \rangle$ can be uniquely expressed as $a(x) + \langle x^N - 1 \rangle$ where $a(x) = \sum_{j=0}^{N-1} a_j x^j \in R[x]$ with $a_0, a_1, \dots, a_{N-1} \in R$. As usual, we will identify $a(x) + \langle x^N - 1 \rangle$ with $a(x)$ for simplicity. Hence

$$R[x]/\langle x^N - 1 \rangle = \{ \sum_{j=0}^{N-1} a_j x^j \mid a_0, a_1, \dots, a_{N-1} \in R \}$$

which is a ring with the operations defined by the usual polynomial operations modulo $x^N - 1$. It is well-known that cyclic codes over R of length N are identified with ideals of the ring $R[x]/\langle x^N - 1 \rangle$, under the identification map $\theta : R^N \rightarrow R[x]/\langle x^N - 1 \rangle$ via $\theta : (a_0, a_1, \dots, a_{N-1}) \mapsto \sum_{j=0}^{N-1} a_j x^j$ ($\forall a_0, a_1, \dots, a_{N-1} \in R$).

There were many literatures on cyclic codes of length N over finite chain rings $R = \mathbb{F}_{p^m}[u]/\langle u^k \rangle$ for various finite fields $\mathbb{F}_{p^m}$ and positive integers k and N . For example: When $k = 2$, Bonnecaze and Udaya [3] investigated cyclic codes and self-dual codes over $\mathbb{F}_2 + u\mathbb{F}_2$ for odd length N . Norton [7] discussed cyclic codes of length N over an arbitrary finite chain ring R systematically when N is not divisible by the characteristic of the residue field $\bar{R}$. Dinh [4] studied cyclic codes and constacyclic codes over Galois extension rings of $\mathbb{F}_2 + u\mathbb{F}_2$ of length $N = 2^s$, and then investigated cyclic codes and constacyclic codes over $\mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$ of length $N = p^s$ in [5].

When $k \geq 3$, for the case of $p = 2$ and $m = 1$ Abualrub and Siap [1] studied cyclic codes over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2$ and $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$ for arbitrary length N , then Al-Ashker and Hamoudeh [2] extended some of the results in [1], and studied cyclic codes of an arbitrary length over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2 + \dots + u^{k-1}\mathbb{Z}_2$ with $u^k = 0$. The rank and minimal spanning set of this family of codes were studied and two open problems were asked: The study of cyclic codes of an arbitrary length over $\mathbb{Z}_p + u\mathbb{Z}_p + u^2\mathbb{Z}_p + \dots + u^{k-1}\mathbb{Z}_p$, where p is a prime integer, $u^k = 0$, and the study of dual and self-dual codes and their properties over these rings. Recently, for the case of $m = 1$ Singh et al. [8] studied cyclic code over the ring $\mathbb{Z}_p[u]/\langle u^k \rangle = \mathbb{Z}_p + u\mathbb{Z}_p + u^2\mathbb{Z}_p + \dots + u^{k-1}\mathbb{Z}_p$ for any prime integer p and positive integer N . A set of generators, the rank and the Hamming distance of these codes were investigated. But for arbitrary prime p and positive integers m, N and $k \geq 3$, the following questions for cyclic codes over R have not been solved completely to the best of our knowledge:

- For each cyclic code $\mathcal{C}$ over R of length N , give a unique representation for $\mathcal{C}$ and provide a clear formula to count the number of codewords in $\mathcal{C}$. Then obtain a formula to count the number of all such cyclic codes.

- Using the representation of any cyclic code $\mathcal{C}$, give the representation of the dual code of $\mathcal{C}$ precisely and determine the self-duality of $\mathcal{C}$.

From now on, let $R = \mathbb{F}_{2^m}[u]/\langle u^k \rangle = \mathbb{F}_{2^m} + u\mathbb{F}_{2^m} + \dots + u^{k-1}\mathbb{F}_{2^m}$ ($u^k = 0$) where $k \geq 2$, and n be an odd positive integer. We focus our attention to consider cyclic codes over R of length $2n$ in this paper.

The present paper is organized as follows. In Section 2, we provide an explicit representation for each cyclic code over R of length $2n$ and give a formula to count the number of codewords in each code. As a corollary, we obtain a formula for the number of all such cyclic codes. In Section 3, we

determine the dual code of each cyclic code over R of length $2n$ and investigate self-duality for such codes. Finally, we list all 293687 cyclic codes and all 791 self-dual cyclic codes of length 14 over $\mathbb{F}_2[u]/\langle u^4 \rangle$ in Section 4.

2 Representation for all distinct cyclic codes over R of length $2n$

In this section, we investigate the structures of $R[x]/\langle x^{2n} - 1 \rangle$ first. Then give an explicit presentation for all distinct cyclic codes over R of length $2n$.

As n is odd, there are pairwise coprime monic irreducible polynomials $f_1(x), \dots, f_r(x)$ in $\mathbb{F}_{2^m}[x]$ such that $x^n - 1 = f_1(x) \dots f_r(x)$. Then we have

$$x^{2n} - 1 = (x^n - 1)^2 = f_1(x)^2 \dots f_r(x)^2.$$

Let $1 \leq j \leq r$. We assume $\deg(f_j(x)) = d_j$ and denote $F_j(x) = \frac{x^{2n}-1}{f_j(x)^2}$. Then $\gcd(F_j(x), f_j(x)^2) = 1$ and there exist $a_j(x), b_j(x) \in \mathbb{F}_{2^m}[x]$ such that

$$a_j(x)F_j(x) + b_j(x)f_j(x)^2 = 1. \quad (1)$$

From now on, we denote $\mathcal{A} = \mathbb{F}_{2^m}[x]/\langle x^{2n} - 1 \rangle$, $\mathcal{K}_j = \mathbb{F}_{2^m}[x]/\langle f_j(x)^2 \rangle$ and set

$$\varepsilon_j(x) \equiv a_j(x)F_j(x) = 1 - b_j(x)f_j(x)^2 \pmod{x^{2n} - 1}. \quad (2)$$

From classical ring theory and coding theory we deduce the following lemma.

Lemma 2.1 *Using the notations above, we have the following:*

(i) $\varepsilon_1(x) + \dots + \varepsilon_r(x) = 1$, $\varepsilon_j(x)^2 = \varepsilon_j(x)$ and $\varepsilon_j(x)\varepsilon_l(x) = 0$ in the ring $\mathcal{A}$ for all $1 \leq j \neq l \leq r$.

(ii) $\mathcal{A} = \mathcal{A}_1 \oplus \dots \oplus \mathcal{A}_r$ where $\mathcal{A}_j = \mathcal{A}\varepsilon_j(x)$ with $\varepsilon_j(x)$ as its multiplicative identity and satisfies $\mathcal{A}_j\mathcal{A}_l = \{0\}$ for all $1 \leq j \neq l \leq r$.

(iii) For any integer j , $1 \leq j \leq r$, define $\varphi_j : a(x) \mapsto \varepsilon_j(x)a(x) \pmod{x^{2n} - 1}$ for any $a(x) \in \mathcal{K}_j$. Then φ_j is a ring isomorphism from $\mathcal{K}_j$ onto $\mathcal{A}_j$ and can be extended to a ring isomorphism from $\mathcal{K}_j[u]/\langle u^k \rangle$ onto $\mathcal{A}_j[u]/\langle u^k \rangle$ in the natural way that: for any $a_0(x), a_1(x), \dots, a_{k-1}(x) \in \mathcal{K}_j$,

$$\varphi_j : \sum_{l=0}^{k-1} a_l(x)u^l \mapsto \sum_{l=0}^{k-1} \varphi_j(a_l(x))u^l = \varepsilon_j(x) \left(\sum_{l=0}^{k-1} a_l(x)u^l \right) \pmod{x^{2n} - 1}.$$

(iv) For any integer j , $1 \leq j \leq r$, $\mathcal{A}_j$ is a cyclic code over $\mathbb{F}_{2^m}$ of length $2n$ with parity check polynomial $f_j(x)^2$.

For any $\alpha(x) = \sum_{i=0}^{2n-1} \alpha_i x^i \in R[x]/\langle x^{2n} - 1 \rangle$ where $\alpha_i = \sum_{j=0}^{k-1} a_{i,j} u^j \in R$ with $a_{i,j} \in \mathbb{F}_{2^m}$ for all $i = 0, 1, \dots, 2n-1$ and $j = 0, 1, \dots, k-1$, we have

$$\alpha(x) = (1, x, \dots, x^{2n-1}) \begin{pmatrix} a_{0,0} & a_{0,1} & \dots & a_{0,k-1} \\ a_{1,0} & a_{1,1} & \dots & a_{1,k-1} \\ \dots & \dots & \dots & \dots \\ a_{2n-1,0} & a_{2n-1,1} & \dots & a_{2n-1,k-1} \end{pmatrix} \begin{pmatrix} 1 \\ u \\ \dots \\ u^{k-1} \end{pmatrix}.$$

Now, define $\Psi(\alpha(x)) = a_0(x) + a_1(x)u + \dots + a_{k-1}(x)u^{k-1} \in \mathcal{A}[u]/\langle u^k \rangle$ where $a_j(x) = a_{0,j} + a_{1,j}x + \dots + a_{2n-1,j}x^{2n-1} \in \mathcal{A}$ for all $j = 0, 1, \dots, k-1$. Then by a direct calculation, one can easily verify the following lemma.

Lemma 2.2 *The map Ψ defined above is a ring isomorphism from $R[x]/\langle x^{2n} - 1 \rangle$ onto $\mathcal{A}[u]/\langle u^k \rangle = \mathcal{A} + u\mathcal{A} + \dots + u^{k-1}\mathcal{A}$ ($u^k = 0$).*

Remark In the rest of this paper, we will identify $R[x]/\langle x^{2n} - 1 \rangle$ with $\mathcal{A}[u]/\langle u^k \rangle$ under the ring isomorphism Ψ .

Therefore, in order to determine cyclic codes over R of length $2n$, i.e., ideals of $R[x]/\langle x^{2n} - 1 \rangle$, it is sufficient to determine ideals of the ring $\mathcal{A}[u]/\langle u^k \rangle$. Precisely, we have the following theorem.

Theorem 2.3 *Using the notations above, the following statements are equivalent:*

- (i) $\mathcal{C}$ is a cyclic code over R of length $2n$.
- (ii) For any integer j , $1 \leq j \leq r$, there is a unique ideal $\mathcal{C}_j$ of the ring $\mathcal{A}_j[u]/\langle u^k \rangle$ such that $\mathcal{C} = \bigoplus_{j=1}^r \mathcal{C}_j$.
- (iii) For any integer j , $1 \leq j \leq r$, there is a unique ideal $\mathcal{C}_j$ of the ring $\mathcal{K}_j[u]/\langle u^k \rangle$ such that $\mathcal{C} = \bigoplus_{j=1}^r \varepsilon_j(x) \mathcal{C}_j \pmod{x^{2n} - 1}$.

Proof (i) $\Leftrightarrow$ (ii). By Lemma 2.1(ii) $\mathcal{A}$ is a direct product of its ideals $\mathcal{A}_1, \dots, \mathcal{A}_r$, which implies $\mathcal{A}[u]/\langle u^k \rangle = \bigoplus_{j=1}^r \mathcal{A}_j[u]/\langle u^k \rangle$. Hence every ideal $\mathcal{C}$ of $\mathcal{A}[u]/\langle u^k \rangle$ is uniquely decomposed as $\mathcal{C} = \bigoplus_{j=1}^r \mathcal{C}_j$, where $\mathcal{C}_j$ is an ideal of $\mathcal{A}_j[u]/\langle u^k \rangle$.

(ii) $\Leftrightarrow$ (iii). Let $1 \leq j \leq r$. By Lemma 2.1(iii), for any ideal $\mathcal{C}_j$ of $\mathcal{A}_j[u]/\langle u^k \rangle$ there is a unique ideal $\mathcal{C}_j$ of $\mathcal{K}_j[u]/\langle u^k \rangle$ such that $\mathcal{C}_j = \varphi_j(\mathcal{C}_j) = \varepsilon_j(x) \mathcal{C}_j$. $\square$

Remark Using the notations of Theorem 2.3, $\mathcal{C} = \bigoplus_{j=1}^r \varepsilon_j(x) \mathcal{C}_j$, where $\mathcal{C}_j$ is an ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$ for $j = 1, \dots, r$, is called a *canonical form decomposition* of the cyclic code $\mathcal{C}$ over R of length $2n$.

By Theorem 2.3, in order to determine ideals of $\mathcal{A}[u]/\langle u^k \rangle$, it is sufficient to determine ideals of $\mathcal{K}_j[u]/\langle u^k \rangle$ for all $j = 1, \dots, r$. Let $1 \leq j \leq r$ and $1 \leq s \leq k$. In the rest of this paper, we adopt the following notations:

- $\mathcal{F}_j = \mathbb{F}_{2^m}[x]/\langle f_j(x) \rangle = \{ \sum_{l=0}^{d_j-1} a_l x^l \mid a_0, a_1, \dots, a_{d_j-1} \in \mathbb{F}_{2^m} \}$.
- $\mathcal{F}_j[u]/\langle u^s \rangle = \{ \sum_{h=0}^{s-1} b_h(x) u^h \mid b_0(x), b_1(x), \dots, b_{s-1}(x) \in \mathcal{F}_j \} = \mathcal{F}_j + u\mathcal{F}_j + \dots + u^{s-1}\mathcal{F}_j$ ($u^s = 0$).

As $f_j(x)$ is a monic irreducible polynomial in $\mathbb{F}_{2^m}[x]$ of degree d_j , the following lemma follows from polynomial theory over finite fields and finite chain ring theory (cf. [7]) immediately. Here, we omit its proof.

Lemma 2.4 *Using the notations above, for any $1 \leq j \leq r$ and $1 \leq s \leq k$ we have the following:*

- (i) $\mathcal{F}_j$ is an extension field of $\mathbb{F}_{2^m}$ with cardinality 2^{md_j} .

(ii) $\mathcal{F}_j[u]/\langle u^s \rangle$ is a finite commutative chain ring with the unique maximal ideal $u(\mathcal{F}_j[u]/\langle u^s \rangle)$, the nilpotency index of u is equal to s and the residue field of $\mathcal{F}_j[u]/\langle u^s \rangle$ is $(\mathcal{F}_j[u]/\langle u^s \rangle)/u(\mathcal{F}_j[u]/\langle u^s \rangle) \cong \mathcal{F}_j$.

(iii) Every element α of $\mathcal{F}_j[u]/\langle u^s \rangle$ has a unique u -adic expansion:

$$\alpha = t_0(x) + ut_1(x) + \dots + u^{s-1}t_{s-1}(x), \quad t_0(x), t_1(x), \dots, t_{s-1}(x) \in \mathcal{F}_j.$$

(iv) The group of invertible elements of $\mathcal{F}_j[u]/\langle u^s \rangle$ is given by

$$(\mathcal{F}_j[u]/\langle u^s \rangle)^\times = \{ \sum_{h=0}^{s-1} b_h(x)u^h \mid b_0(x) \neq 0, b_0(x), b_1(x), \dots, b_{s-1}(x) \in \mathcal{F}_j \}$$

and $|(\mathcal{F}_j[u]/\langle u^s \rangle)^\times| = (2^{md_j} - 1)2^{(s-1)md_j}$. Then for any nonzero element α of $\mathcal{F}_j[u]/\langle u^s \rangle$ there is a unique integer e , $0 \leq e \leq s-1$, such that $\alpha = u^e \omega$ for some $\omega \in (\mathcal{F}_j[u]/\langle u^s \rangle)^\times$.

(v) Denote $\mathcal{T}_j = \{ \sum_{l=0}^{d_j-1} a_l x^l \mid a_0, a_1, \dots, a_{d_j-1} \in \mathbb{F}_{2^m} \} \subset \mathcal{K}_j$. Then the subset $\mathcal{T}_j$ of $\mathcal{K}_j$ satisfies the following properties:

◇ For any $a(x) \in \mathcal{K}_j$, there is a unique $t(x) \in \mathcal{T}_j$ such that $a(x) \equiv t(x) \pmod{f_j(x)}$ as polynomials in $\mathbb{F}_{2^m}[x]$;

◇ $t_1(x) \not\equiv t_2(x) \pmod{f_j(x)}$, for any $t_1(x), t_2(x) \in \mathcal{T}_j$ satisfying $t_1(x) \neq t_2(x)$.

Furthermore, $\mathcal{K}_j$ is a finite chain ring with the maximal ideal $f_j(x)\mathcal{K}_j$ where $f_j(x)^2 = 0$, and every element of $\mathcal{K}_j$ has a unique $f_j(x)$ -adic expansion:

$$t_0(x) + f_j(x)t_1(x), \quad t_0(x), t_1(x) \in \mathcal{T}_j.$$

(vi) Every element ϑ of $\mathcal{K}_j[u]/\langle u^s \rangle$ can be uniquely expressed as

$$\vartheta = \alpha_0 + f_j(x)\alpha_1, \quad \alpha_0, \alpha_1 \in \mathcal{T}_j[u]/\langle u^s \rangle,$$

where $\mathcal{T}_j[u]/\langle u^s \rangle = \{ \sum_{l=0}^{s-1} t_l(x)u^l \mid t_l(x) \in \mathcal{T}_j, 0 \leq l \leq s-1 \} \subset \mathcal{K}_j[u]/\langle u^s \rangle$.

Remark $\mathcal{F}_j = \{ \sum_{l=0}^{d_j-1} a_l x^l \mid a_0, a_1, \dots, a_{d_j-1} \in \mathbb{F}_{2^m} \}$ which is a finite field with operations defined by the usual polynomial operations modulo $f_j(x)$. But $\mathcal{T}_j$ is only a subset of $\mathcal{K}_j$ since the operations on $\mathcal{K}_j$ are defined by the usual polynomial operations modulo $f_j(x)^2$. In spite of this, we have $\mathcal{F}_j = \mathcal{T}_j$ as a set. Therefore, $\mathcal{F}_j[u]/\langle u^s \rangle = \{ \sum_{i=0}^{s-1} \xi_i u^i \mid \xi_0, \xi_1, \dots, \xi_{s-1} \in \mathcal{F}_j \} = \mathcal{T}_j[u]/\langle u^s \rangle$ as a set. In this sense, $\mathcal{F}_j[u]/\langle u^s \rangle$ is not a subring but a subset of $\mathcal{K}_j[u]/\langle u^s \rangle$.

Now, define a map $\tau : a(x) \mapsto a(x) \pmod{f_j(x)}$ for all $a(x) \in \mathcal{K}_j = \mathbb{F}_{2^m}[x]/\langle f_j(x)^2 \rangle$. It is clear that τ is a surjective ring homomorphism from $\mathcal{K}_j$ onto $\mathcal{F}_j = \mathbb{F}_{2^m}[x]/\langle f_j(x) \rangle$ with kernel $\text{Ker}(\tau) = f_j(x)\mathcal{K}_j$. Then we extend τ to $\mathcal{K}_j[u]/\langle u^s \rangle = \mathcal{K}_j + u\mathcal{K}_j + \dots + u^{s-1}\mathcal{K}_j$ ($u^s = 0$) by

$$\tau : \sum_{l=0}^{s-1} a_l(x)u^l \mapsto \sum_{l=0}^{s-1} \tau(a_l(x))u^l \quad (\forall a_0(x), a_1(x), \dots, a_{s-1}(x) \in \mathcal{K}_j).$$

Then τ is a surjective ring homomorphism from $\mathcal{K}_j[u]/\langle u^s \rangle$ onto $\mathcal{F}_j[u]/\langle u^s \rangle$ with kernel $\text{Ker}(\tau) = f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle) = \{ f_j(x)\xi \mid \xi \in \mathcal{T}_j[u]/\langle u^s \rangle \}$.

Let $\alpha \in \mathcal{F}_j[u]/\langle u^s \rangle$ and $f_j(x)\xi \in f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$ where $\xi \in \mathcal{T}_j[u]/\langle u^s \rangle$. As $\mathcal{T}_j[u]/\langle u^s \rangle = \mathcal{F}_j[u]/\langle u^s \rangle$ as a set, we regard ξ as an element of $\mathcal{F}_j[u]/\langle u^s \rangle$ and calculate the product $\xi\alpha$ of ξ and α in the ring $\mathcal{F}_j[u]/\langle u^s \rangle$. Also, we can regard $\xi\alpha$ as an element of $\mathcal{T}_j[u]/\langle u^s \rangle$. Then by Lemma 2.4(vi), we obtain a unique element $f_j(x)(\xi\alpha)$ in $f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$. Now, we define

$$\alpha \cdot f_j(x)\xi = f_j(x)(\xi\alpha) \in f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle).$$

Especially, we have $\alpha \cdot f_j(x) = f_j(x)\alpha \in f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$ if $\xi = 1$. Then $f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$ is an $\mathcal{F}_j[u]/\langle u^s \rangle$ -module. Moreover, we have

Lemma 2.5 *With the scalar multiplication defined above, the map ψ_j defined by $\psi_j(\alpha) = f_j(x)\alpha \in \mathcal{K}_j[u]/\langle u^s \rangle$ ($\forall \alpha \in \mathcal{F}_j[u]/\langle u^s \rangle$) is an $\mathcal{F}_j[u]/\langle u^s \rangle$ -module isomorphism from $\mathcal{F}_j[u]/\langle u^s \rangle$ onto $f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$.*

Proof By Lemma 2.4(vi), every element ϑ of $\mathcal{K}_j[u]/\langle u^s \rangle$ can be uniquely expressed as $\vartheta = \alpha_0 + f_j(x)\alpha_1$ where $\alpha_0, \alpha_1 \in \mathcal{T}_j[u]/\langle u^s \rangle = \mathcal{F}_j[u]/\langle u^s \rangle$. From this and by $f_j(x)^2 = 0$ in $\mathcal{K}_j[u]/\langle u^s \rangle$, we deduce that $f_j(x)\vartheta = f_j(x)\alpha_0$. Hence ψ_j is a bijection. Moreover, for any $\alpha_1, \alpha_2 \in \mathcal{F}_j[u]/\langle u^s \rangle$ we assume that $\alpha_1 + \alpha_2 = \beta$ and $\alpha_1\alpha_2 = \gamma$ in $\mathcal{F}_j[u]/\langle u^s \rangle$. Now, we regard α_1 and α_2 as elements of $\mathcal{K}_j[u]/\langle u^s \rangle$ and assume $\alpha_1 + \alpha_2 = \zeta$ and $\alpha_1\alpha_2 = \eta$ in $\mathcal{K}_j[u]/\langle u^s \rangle$. Since τ is a surjective ring homomorphism from $\mathcal{K}_j[u]/\langle u^s \rangle$ onto $\mathcal{F}_j[u]/\langle u^s \rangle$, by the definition of τ we have $\tau(\zeta) = \alpha_1 + \alpha_2 = \beta$ and $\tau(\eta) = \alpha_1\alpha_2 = \gamma$, which imply $\zeta = \beta + f_j(x)\beta_1$ and $\eta = \gamma + f_j(x)\gamma_1$ for some $\beta_1, \gamma_1 \in \mathcal{T}_j[u]/\langle u^s \rangle$. By $f_j(x)^2 = 0$ in $\mathcal{K}_j[u]/\langle u^s \rangle$, we have

$$\begin{aligned} \psi_j(\alpha_1 + \alpha_2) &= f_j(x)\beta = f_j(x)(\beta + f_j(x)\beta_1) = f_j(x)\zeta = f_j(x)(\alpha_1 + \alpha_2) \\ &= f_j(x)\alpha_1 + f_j(x)\alpha_2 = \psi_j(\alpha_1) + \psi_j(\alpha_2) \end{aligned}$$

and $\psi_j(\alpha_1\alpha_2) = f_j(x)\gamma = f_j(x)(\gamma + f_j(x)\gamma_1) = f_j(x)\eta = f_j(x)(\alpha_1\alpha_2) = \alpha_1(f_j(x)\alpha_2) = \alpha_1\psi_j(\alpha_2)$. Hence ψ_j is an $\mathcal{F}_j[u]/\langle u^s \rangle$ -module isomorphism from $\mathcal{F}_j[u]/\langle u^s \rangle$ onto $f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$. $\square$

Remark By Lemma 2.5, every element of $f_j(x)(\mathcal{K}_j[u]/\langle u^s \rangle)$ can be uniquely expressed as: $f_j(x)\alpha = \psi_j(\alpha)$, $\alpha \in \mathcal{F}_j[u]/\langle u^s \rangle$, where we regard α in the expression $f_j(x)\alpha$ as an element of $\mathcal{T}_j[u]/\langle u^s \rangle$ in the rest of this paper.

Then we present all distinct ideals of the $\mathcal{K}_j[u]/\langle u^k \rangle$.

Theorem 2.6 *Using the notations above, all distinct ideals of $\mathcal{K}_j[u]/\langle u^k \rangle$ are given by the following table:*

case	number of ideals	C_j (ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$)	$ C_j $
I.	$k + 1$	$\bullet \langle u^i \rangle \quad (0 \leq i \leq k)$	$2^{2md_j(k-i)}$
II.	k	$\bullet \langle u^s f_j(x) \rangle \quad (0 \leq s \leq k - 1)$	$2^{md_j(k-s)}$
III.	$\Omega_1(2^{md_j}, k)$	$\bullet \langle u^i + u^t f_j(x)\omega \rangle$ $(\omega \in (\mathcal{F}_j[u]/\langle u^{i-t} \rangle)^\times, t \geq 2i - k,$ $0 \leq t < i \leq k - 1)$	$2^{2md_j(k-i)}$
IV.	$\Omega_2(2^{md_j}, k)$	$\bullet \langle u^i + u^t f_j(x)\omega \rangle$ $(\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times, t < 2i - k,$ $0 \leq t < i \leq k - 1)$	$2^{md_j(k-t)}$
V.	$\frac{1}{2}k(k - 1)$	$\bullet \langle u^i, u^s f_j(x) \rangle$ $(0 \leq s < i \leq k - 1)$	$2^{md_j(2k-(i+s))}$
VI.	$(2^{md_j} - 1) \cdot \Gamma(2^{md_j}, k)$	$\bullet \langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$ $(\omega \in (\mathcal{F}_j[u]/\langle u^{s-t} \rangle)^\times, i + s \leq k + t - 1,$ $0 \leq t < s < i \leq k - 1)$	$2^{md_j(2k-(i+s))}$

where

$$\diamond \Omega_1(2^{md_j}, k) = \begin{cases} \frac{2^{md_j(\frac{k}{2}+1)} + 2^{md_j \cdot \frac{k}{2}} - 2}{2^{md_j-1}} - (k+1), & \text{if } k \text{ is even;} \\ \frac{2(2^{md_j \cdot \frac{k+1}{2}} - 1)}{2^{md_j-1}} - (k+1), & \text{if } k \text{ is odd.} \end{cases}$$

$$\diamond \Omega_2(2^{md_j}, k) = \begin{cases} (2^{md_j} - 1) \sum_{i=\frac{k}{2}+1}^{k-1} (2i - k) 2^{md_j(k-i-1)}, & \text{if } k \text{ is even;} \\ (2^{md_j} - 1) \sum_{i=\frac{k+1}{2}}^{k-1} (2i - k) 2^{md_j(k-i-1)}, & \text{if } k \text{ is odd.} \end{cases}$$

$\diamond \Gamma(2^{md_j}, k)$ can be calculated by the following recurrence formula:

$$\Gamma(2^{md_j}, \rho) = 0 \text{ for } \rho = 1, 2, 3, \Gamma(2^{md_j}, \rho) = 1 \text{ for } \rho = 4;$$

$$\Gamma(2^{md_j}, \rho) = \Gamma(2^{md_j}, \rho - 1) + \sum_{s=1}^{\lfloor \frac{\rho}{2} \rfloor - 1} (\rho - 2s - 1) 2^{md_j(s-1)} \text{ for } \rho \geq 5.$$

Therefore, the number of all distinct ideals of the ring $\mathcal{K}_j[u]/\langle u^k \rangle$ is equal to

$$N_{(2^m, d_j, k)} = 1 + \frac{k(k+3)}{2} + \Omega_1(2^{md_j}, k) + \Omega_2(2^{md_j}, k) + (2^{md_j} - 1)\Gamma(2^{md_j}, k).$$

Proof In order to simplify the notations, we denote $\mathcal{Y}_j = \mathcal{K}_j[u]/\langle u^k \rangle$ and $\mathcal{R}_j = \mathcal{F}_j[u]/\langle u^k \rangle$. By Lemma 2.4(ii), $\mathcal{R}_j$ is a finite chain ring with maximal ideal $u\mathcal{R}_j$, the nilpotency index of u is equal to k and the residue field of $\mathcal{R}_j$ is $\mathcal{R}_j/(u\mathcal{R}_j) \cong \mathcal{F}_j$. Then from finite chain ring theory (cf. [7]), we deduce that all ideals of $\mathcal{R}_j$ is given by $u^l\mathcal{R}_j$, where $0 \leq l \leq k$, and $|u^l\mathcal{R}_j| = |\mathcal{F}_j|^{k-l} = 2^{md_j(k-l)}$. Moreover, we have $f_j(x)\mathcal{Y}_j = f_j(x)\mathcal{R}_j$ by Lemma 2.5.

Let C be any ideal of $\mathcal{Y}_j$. We define

$$\text{Tor}_0(C) = \tau(C), \text{Tor}_1(C) = \tau(\{\alpha \in \mathcal{Y}_j \mid f_j(x)\alpha \in C\}).$$

Then it can be easily verify that both $\text{Tor}_0(C)$ and $\text{Tor}_1(C)$ are ideals of $\mathcal{R}_j$ satisfying $\text{Tor}_0(C) \subseteq \text{Tor}_1(C)$. Hence there is a unique pair (i, s) of integers, $0 \leq s \leq i \leq k$, such that $\text{Tor}_0(C) = u^i\mathcal{R}_j$ and $\text{Tor}_1(C) = u^s\mathcal{R}_j$.

Let $\tau|_C$ be the restriction of τ to C . Then $\tau|_C$ is a surjective ring homomorphism from C onto $\text{Tor}_0(C)$ with kernel $\text{Ker}(\tau|_C) = f_j(x)\text{Tor}_1(C)$. As $|\text{Ker}(\tau|_C)| = |\text{Tor}_1(C)|$, by the ring isomorphism theorems it follows that

$$|C| = |\text{Tor}_0(C)||\text{Tor}_1(C)| = |u^i\mathcal{R}_j||u^s\mathcal{R}_j| = 2^{md_j(2k-(i+s))}. \quad (3)$$

Then we have the following cases.

Case (i) $s = i$. In this case, we have $|C| = 2^{2md_j(k-i)}$ by Equation (3).

When $s = k$, then $i = k$ and $C = \{0\} = u^k\mathcal{Y}_j = \langle u^k \rangle$.

Let $0 \leq i \leq k - 1$. By $u^i \in u^i\mathcal{R}_j = \text{Tor}_0(C)$, there exists $\alpha \in \mathcal{Y}_j$ such that $u^i + f_j(x)\alpha \in C$. Then by Lemma 2.5 we can assume that $\alpha \in \mathcal{R}_j$.

It is obvious that $\langle u^i + f_j(x)\alpha \rangle = \mathcal{Y}_j(u^i + f_j(x)\alpha) \subseteq C$. Conversely, let $\xi \in C$. By $\text{Tor}_0(C) = u^i\mathcal{R}_j$ and Lemma 2.5 we have that $\xi = u^i\beta + f_j(x)\gamma$ for some $\beta, \gamma \in \mathcal{R}_j$, which implies $f_j(x)(\gamma - \alpha\beta) = \xi - (u^i + f_j(x)\alpha)\beta \in C$. By Lemma 2.5 we may regard $\gamma - \alpha\beta$ as an element of $\mathcal{R}_j$. Hence $\gamma - \alpha\beta = \tau(\gamma - \alpha\beta) \in \text{Tor}_1(C) = u^s\mathcal{R}_j$, which implies that $\gamma - \alpha\beta = u^i\delta$ for some $\delta \in \mathcal{R}_j$. Therefore, we have $\xi = (u^i + f_j(x)\alpha)\beta + (u^i + f_j(x)\alpha)f_j(x)\delta = (u^i + f_j(x)\alpha)(\beta + f_j(x)\delta) \in \langle u^i + f_j(x)\alpha \rangle$. Hence $C = \langle u^i + f_j(x)\alpha \rangle$.

When $i = 0$, $1 + f_j(x)\alpha \in C$. By $(1 + f_j(x)\alpha)(1 - f_j(x)\alpha) = 1 - f_j(x)^2\alpha^2 = 1$ in $\mathcal{Y}_j$, we see that $1 + f_j(x)\alpha$ is an invertible element of $\mathcal{Y}_j$, which implies $C = \mathcal{Y}_j = \langle u^0 \rangle$. In the following, we assume $1 \leq i \leq k - 1$.

Obviously, we have $f_j(x)u^i = f_j(x)(u^i + f_j(x)\alpha) \in C$. By Lemma 2.4(iv), we can write α as $\alpha = u^i\beta$ or $\alpha = u^t\omega + u^i\beta$ for some $\beta \in \mathcal{R}_j$, $0 \leq t \leq i - 1$ and $\omega \in \mathcal{R}_j^\times$.

When $\alpha = u^i\beta$, we have $u^i = u^i + f_j(x)\alpha - (f_j(x)u^i)\beta \in C$, and so $C = \langle u^i \rangle$.

Let $\alpha = u^t\omega + u^i\beta$. Then we have $u^i + u^t f_j(x)\omega = u^i + f_j(x)\alpha - (f_j(x)u^i)\beta \in C$, and hence $C = \langle u^i + u^t f_j(x)\omega \rangle$. In this case, we have $u^{k-i+t} f_j(x) = u^{k-i}\omega^{-1}(u^i + u^t f_j(x)\omega) \in C$ where ω^{-1} is the inverse of ω in $\mathcal{R}_j$. Therefore, $u^{k-i+t} \in \text{Tor}_1(C) = u^i \mathcal{R}_j$, which implies $k - i + t \geq i$, and hence $t \geq 2i - k$.

Now, let $\omega_1, \omega_2 \in \mathcal{R}_j^\times$ and $0 \leq t_1, t_2 \leq k - 1$ satisfying $C = \langle u^i + u^{t_1} f_j(x)\omega_1 \rangle = \langle u^i + u^{t_2} f_j(x)\omega_2 \rangle$. Then $f_j(x)(u^{t_1}\omega_1 - u^{t_2}\omega_2) = (u^i + u^{t_1} f_j(x)\omega_1) - (u^i + u^{t_2} f_j(x)\omega_2) \in C$, which implies $u^{t_1}\omega_1 - u^{t_2}\omega_2 \in \text{Tor}_1(C) = u^i \mathcal{R}_j$. From this we deduce that $t_1 = t_2 = t$ and $u^t(\omega_1 - \omega_2) \in u^i \mathcal{R}_j$. Then by finite chain ring theory (cf. [7]) and $u^t(\omega_1 - \omega_2) \in u^i \mathcal{R}_j$, it follows that $\omega_1 \equiv \omega_2 \pmod{u^{i-t}}$, i.e., $\omega_1 = \omega_2$ as elements of $(\mathcal{R}_j / (u^{i-t} \mathcal{R}_j))^\times$. Finally, by $\mathcal{R}_j = \mathcal{F}_j[u] / \langle u^k \rangle$ we have $(\mathcal{R}_j / (u^{i-t} \mathcal{R}_j))^\times = \mathcal{F}_j[u] / \langle u^{i-t} \rangle$ up to a natural ring isomorphism.

As stated above, we conclude that all distinct ideals C of $\mathcal{Y}_j$ is given by (I) and (III) in the table.

Case (ii) $i = k$ and $0 \leq s \leq k - 1$.

In this case, we have $\text{Tor}_0(C) = \{0\}$ and $|C| = 2^{md_j(2k-(k+s))} = 2^{md_j(k-s)}$ by Equation (3). Moreover, by $\text{Tor}_1(C) = u^s \mathcal{R}_j$ it can be easily verified that all distinct ideals of $\mathcal{Y}_j$ are given by: (II) $C = \langle u^s f_j(x) \rangle$ where $0 \leq s \leq k - 1$.

Case (iii) $s = 0$ and $1 \leq i \leq k - 1$.

In this case, we have $|C| = 2^{md_j(2k-i)}$ by Equation (3). Moreover, by $1 \in \text{Tor}_1(C) = u^0 \mathcal{R}_j$ we conclude that $f_j(x) = u^0 f_j(x) \in C$. Then by $\text{Tor}_0(C) = u^i \mathcal{R}_j$ it follows that $C = \langle u^i, f_j(x) \rangle$ immediately.

Case (iv) $1 \leq s < i \leq k - 1$. In this case, $|C| = 2^{md_j(2k-(i+s))}$ by (3).

By $\text{Tor}_0(C) = u^i \mathcal{R}_j$ and $\text{Tor}_1(C) = u^s \mathcal{R}_j$ we have $u^s f_j(x) \in C$ and there exists $\alpha \in \mathcal{R}_j$ such that $u^i + f_j(x)\alpha \in C$, which implies $\langle u^i + f_j(x)\alpha, u^s f_j(x) \rangle \subseteq C$. Conversely, let $\xi \in C$. By $\text{Tor}_0(C) = u^i \mathcal{R}_j$ there exist $\beta, \gamma \in \mathcal{R}_j$ such that $\xi = u^i\beta + f_j(x)\gamma$. Then from $f_j(x)(\gamma - \alpha\beta) = \xi - (u^i + f_j(x)\alpha)\beta \in C$ where $\gamma - \alpha\beta \in \mathcal{R}_j$, we deduce $\gamma - \alpha\beta \in \text{Tor}_1(C) = u^s \mathcal{R}_j$, which implies $\gamma - \alpha\beta = u^s\delta$ for some $\delta \in \mathcal{R}_j$, and so $\xi = (u^i + f_j(x)\alpha)\beta + (f_j(x)u^s)\delta \in \langle u^i + f_j(x)\alpha, f_j(x)u^s \rangle$. Hence $C = \langle u^i + f_j(x)\alpha, u^s f_j(x) \rangle$. Furthermore, by $u^s f_j(x) \in C$ and an argument similar to the proof of Case (i) we can assume that $\alpha = 0$ or $\alpha = u^t\omega$, where $\omega \in \mathcal{R}_j^\times$ and $0 \leq t \leq s - 1$.

(iv-1) When $\alpha = 0$, we have $C = \langle u^i, u^s f_j(x) \rangle$ given by (V) in the table.

(iv-2) Let $C = \langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$ where $\omega \in \mathcal{R}_j^\times$ and $0 \leq t \leq s - 1$.

Assume $C = \langle u^i + u^{t_1} f_j(x)\omega_1, u^s f_j(x) \rangle = \langle u^i + u^{t_2} f_j(x)\omega_2, u^s f_j(x) \rangle$ where $0 \leq t_1, t_2 \leq s - 1$ and $\omega_1, \omega_2 \in \mathcal{R}_j^\times$. Then $f_j(x)(u^{t_1}\omega_1 - u^{t_2}\omega_2) = (u^i + u^{t_1} f_j(x)\omega_1) - (u^i + u^{t_2} f_j(x)\omega_2) \in C$, which implies $u^{t_1}\omega_1 - u^{t_2}\omega_2 \in \text{Tor}_1(C) = u^s \mathcal{R}_j$. From this and by an argument similar to the proof of Case (i), we

deduce that $t_1 = t_2 = t$ and $\omega_1 \equiv \omega_2 \pmod{u^{s-t}\mathcal{R}_j}$. By the latter condition, we have $\omega = \omega_1 = \omega_2 \in (\mathcal{R}_j/(u^{s-t}\mathcal{R}_j))^\times$. From this and by $\mathcal{R}_j/(u^{s-t}\mathcal{R}_j) = \mathcal{F}_j[u]/\langle u^{s-t} \rangle$, we deduce that all distinct ideals of $\mathcal{Y}_j$ are given by: $C = \langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$ where $\omega \in (\mathcal{F}_j[u]/\langle u^{s-t} \rangle)^\times$ in this case.

As $f_j(x)u^{k-i+t} = u^{k-i}\omega^{-1}(u^i + u^t f_j(x)\omega) \in C$, we have $u^{k-i+t} \in \text{Tor}_1(C) = u^s \mathcal{R}_j$, which implies $k - i + t \geq s$. So we have one of the following two cases:

($\diamond$ -1) $k - i + t = s$, i.e., $s - t = k - i$.

In this case, by $f_j(x)u^s = u^{k-i}\omega^{-1}(u^i + u^t f_j(x)\omega)$ we have $C = \langle u^i + u^t f_j(x)\omega \rangle$ and $2i > i + s = k + t$, i.e., $t < 2i - k$. Furthermore, we have $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$ and $|C| = 2^{md_j(2k-(i+s))} = 2^{md_j(2k-(k+t))} = 2^{md_j(k-t)}$. Hence C is given by (IV) in the table.

($\diamond$ -2) $k - i + t > s$, i.e., $i + s \leq k + t - 1$. In this case, C is given by (VI) in the table.

As stated above, we conclude that all distinct ideals and the number of elements in each ideal of $\mathcal{Y}_j$ are given by (I)–(VI) of the table.

It is obvious that the number of ideals in (I), (II) and (V) is equal to $k + 1$, k and $\frac{1}{2}k(k - 1)$ respectively. Then we count the number of ideals in (III), (IV) and (V) respectively. In order to simplify the notation, we denote $q_j = 2^{md_j}$ in the following. Then by Lemma 2.4(iv) we know that $(\mathcal{F}_j[u]/\langle u^l \rangle)^\times = (q_j - 1)q_j^{l-1}$ for all $l = 1, \dots, k$.

First, we count the number of ideals in (III). Let $0 \leq t < i \leq k - 1$. Then we have one of the following two cases:

($\diamond$ -1) k is even. In this case, $t \geq 2i - k$ if and only if i and t satisfy one of the following two conditions:

($\diamond$ -1-1) $i \leq \frac{k}{2}$, i.e., $2i \leq k$, and $0 \leq t \leq i - 1$. In this case, the number of

ideals is equal to $N_1 = \sum_{i=1}^{\frac{k}{2}} \sum_{t=0}^{i-1} (q_j - 1)q_j^{i-t-1} = \frac{q_j^{\frac{k}{2}+1} - 1}{q_j - 1} - (\frac{k}{2} + 1)$.

($\diamond$ -1-2) $i \geq \frac{k}{2} + 1$, i.e., $2i > k$, and $2i - k \leq t \leq i - 1$. In this case, the number of ideals is equal to $N_2 = \sum_{i=\frac{k}{2}+1}^{k-1} \sum_{t=2i-k}^{i-1} (q_j - 1)q_j^{i-t-1} = \frac{q_j^{\frac{k}{2}-1} - 1}{q_j - 1} - \frac{k}{2}$.

Therefore, the number of ideals in (III) is equal to $\Omega_1(q_j, k) = N_1 + N_2 = \frac{q_j^{\frac{k}{2}+1} + q_j^{\frac{k}{2}-2} - 2}{q_j - 1} - (k + 1)$, where k is even.

($\diamond$ -2) k is odd. In this case, $t \geq 2i - k$ if and only if i and t satisfy one of the following two conditions:

($\diamond$ -2-1) $i \leq \frac{k-1}{2}$, i.e., $2i \leq k$, and $0 \leq t \leq i - 1$. Then the number of ideals

is equal to $N_1 = \sum_{i=1}^{\frac{k-1}{2}} \sum_{t=0}^{i-1} (q_j - 1)q_j^{i-t-1} = \frac{q_j^{\frac{k+1}{2}} - 1}{q_j - 1} - \frac{k+1}{2}$.

($\diamond$ -2-2) $i \geq \frac{k-1}{2} + 1$, i.e., $2i > k$, and $2i - k \leq t \leq i - 1$. Then the number of ideals is equal to $N_2 = \sum_{i=\frac{k-1}{2}+1}^{k-1} \sum_{t=2i-k}^{i-1} (q_j - 1)q_j^{i-t-1} = \frac{q_j^{\frac{k+1}{2}} - 1}{q_j - 1} - \frac{k+1}{2}$.

Therefore, the number of ideals in (III) is equal to $\Omega_1(q_j, k) = N_1 + N_2 = \frac{2q_j^{\frac{k+1}{2}} - 2}{q_j - 1} - (k + 1)$, where k is odd.

Next we count the number of ideals in (IV). Let $0 \leq t < i \leq k - 1$. Then we have one of the following two cases:

($\diamond$ -1) k is even. Then $t < 2i - k$ if and only if $2i > k$, i.e., $i \geq \frac{k}{2} + 1$, and $0 \leq t \leq 2i - k - 1$. Hence the number of ideals in (IV) is equal to $\Omega_2(q_j, k) = \sum_{i=\frac{k}{2}+1}^{k-1} \sum_{t=0}^{2i-k-1} (q_j - 1)q_j^{k-i-1} = (q_j - 1) \sum_{i=\frac{k}{2}+1}^{k-1} (2i - k)q_j^{k-i-1}$.

($\diamond$ -2) k is odd. Then $t < 2i - k$ if and only if $2i > k$, i.e., $i \geq \frac{k+1}{2}$, and $0 \leq t \leq 2i - k - 1$. Hence the number of ideals in (IV) is equal to $\Omega_2(q_j, k) = \sum_{i=\frac{k+1}{2}}^{k-1} \sum_{t=0}^{2i-k-1} (q_j - 1)q_j^{k-i-1} = (q_j - 1) \sum_{i=\frac{k+1}{2}}^{k-1} (2i - k)q_j^{k-i-1}$.

Finally, denote $\Gamma(q_j, k) = \sum_{t=0}^{k-4} \sum_{t+1 \leq s \leq \frac{k+t}{2}-1} \sum_{s+1 \leq i \leq k+t-1-s} q_j^{s-t-1} + \sum_{1 \leq s \leq \frac{k}{2}-1} \sum_{s+1 \leq i \leq k-1-s} q_j^{s-1}$. Then the number of ideals in (VI) is equal to $(q_j - 1)\Gamma(q_j, k)$, where

$$\begin{aligned} \Gamma(q_j, k) &= \sum_{t=1}^{k-4} \sum_{t+1 \leq s \leq \frac{k+t}{2}-1} \sum_{s+1 \leq i \leq k+t-1-s} q_j^{s-t-1} + \sum_{1 \leq s \leq \frac{k}{2}-1} \sum_{s+1 \leq i \leq k-1-s} q_j^{s-1} \\ &= \sum_{t'=0}^{(k-1)-4} \sum_{t'+1 \leq s' \leq \frac{(k-1)+t'}{2}-1} \sum_{s'+1 \leq i' \leq (k-1)+t'-1-s'} q_j^{s'-t'-1} \\ &\quad + \sum_{s=1}^{\lfloor \frac{k}{2} \rfloor - 1} (k - 2s - 1)q_j^{s-1} \\ &= \Gamma(q_j, k-1) + \sum_{s=1}^{\lfloor \frac{k}{2} \rfloor - 1} (k - 2s - 1)q_j^{s-1} \end{aligned}$$

when $k \geq 5$.

If $1 \leq k \leq 3$, there is no triple (t, s, i) of integers satisfying $0 \leq t < s < i \leq k-1$ and $i + s \leq k + t - 1$. In this case, the number of ideals in (VI) is equal to 0. Then we set $\Gamma(q_j, k) = 0$ for $k = 1, 2, 3$.

If $k = 4$, there is a unique triple $(t, s, i) = (0, 1, 2)$ of integers satisfying $0 \leq t < s < i \leq k-1$ and $i + s \leq k + t - 1$. In this case, all distinct ideals in (VI) are given by $\langle u^2 + f_j(x)\omega, uf_j(x) \rangle$, where $\omega \in (\mathcal{F}_j[u]/\langle u \rangle)^\times = \mathcal{F}_j^\times$ and $|\mathcal{F}_j^\times| = q_j - 1$. Then we set $\Gamma(q_j, 4) = 1$.

Therefore, the number of ideals of $\mathcal{Y}_j$ is equal to $k + 1 + k + \Omega_1(q_j, k) + \Omega_2(q_j, k) + \frac{1}{2}k(k-1) + (q_j - 1)\Gamma(q_j, k) = 1 + \frac{1}{2}k(k+3) + \Omega_1(q_j, k) + \Omega_2(q_j, k) + (q_j - 1)\Gamma(q_j, k)$. $\square$

Corollary 2.7 *Using the notations in Theorems 2.3 and 2.6, the number of cyclic codes over $\mathbb{F}_{2^m}[u]/\langle u^k \rangle$ of length $2n$ is equal to $\prod_{j=1}^r N_{(2^m, d_j, k)}$.*

Proof It follows from Theorems 2.3 and 2.6 immediately. $\square$

3 Dual codes and self-duality of cyclic codes over R of length $2n$

In this section, we give the dual code of each cyclic code over the ring $R = \mathbb{F}_{2^m}[u]/\langle u^k \rangle$ of length $2n$ where n is odd, and then investigate the self-duality of these codes.

For any $\alpha = (\alpha_0, \alpha_1, \dots, \alpha_{2n-1}), \beta = (\beta_0, \beta_1, \dots, \beta_{2n-1}) \in R^{2n}$, where $\alpha_j, \beta_j \in R$ and $0 \leq j \leq 2n-1$, recall that the usual *Euclidian inner product* of α and β are defined by $[\alpha, \beta]_E = \sum_{j=0}^{2n-1} \alpha_j \beta_j \in R$. It is known that $[-, -]_E$ is a symmetric and non-degenerate bilinear form on the R -module R^{2n} . Let C be a linear code over R of length $2n$, i.e., an R -submodule of R^{2n} . The *Euclidian dual code* of C is defined by $C^{\perp_E} = \{\alpha \in R^{2n} \mid [\alpha, \beta]_E = 0, \forall \beta \in C\}$, and C is said to be *self-dual* if $C = C^{\perp_E}$. As usual, we will identify $\alpha \in R^{2n}$ with $\alpha(x) = \sum_{j=0}^{2n-1} \alpha_j x^j \in R[x]/\langle x^{2n} - 1 \rangle$ in this paper. Then we define

$$\varrho(\alpha(x)) = \alpha(x^{-1}) = \alpha_0 + \sum_{j=1}^{2n-1} \alpha_j x^{2n-j}, \quad \forall \alpha(x) \in R[x]/\langle x^{2n} - 1 \rangle.$$

It is clear that ϱ is a ring automorphism of $R[x]/\langle x^{2n} - 1 \rangle$ satisfying $\varrho^{-1} = \varrho$. Now, by a direct calculation we get the following lemma.

Lemma 3.1 *Let $\alpha, \beta \in R^{2n}$. Then $[\alpha, \beta]_E = 0$ if $\alpha(x)\varrho(\beta(x)) = 0$ in the ring $R[x]/\langle x^{2n} - 1 \rangle$.*

By Lemma 2.2 and the remark followed this lemma, we have $R[x]/\langle x^{2n} - 1 \rangle = \mathcal{A}[u]/\langle u^k \rangle$ where $\mathcal{A} = \mathbb{F}_{2^m}[x]/\langle x^{2n} - 1 \rangle$. It is obvious that the restriction of ϱ to $\mathcal{A}$ is a ring automorphism of $\mathcal{A}$. We still denote this automorphism by ϱ , i.e., $\varrho(a(x)) = a(x^{-1})$ for any $a(x) \in \mathcal{A}$.

Let $1 \leq j \leq r$. By Equations (1) and (2) in Section 2, we have

$$\varrho(\varepsilon_j(x)) = a_j(x^{-1})F_j(x^{-1}) = 1 - b_j(x^{-1})f_j(x^{-1})^2 \text{ in } \mathcal{A}. \quad (4)$$

For any polynomial $f(x) = \sum_{l=0}^d a_l x^l \in \mathbb{F}_{2^m}[x]$ of degree $d \geq 1$, recall that the *reciprocal polynomial* of $f(x)$ is defined as $\tilde{f}(x) = x^d f(\frac{1}{x}) = \sum_{l=0}^d a_l x^{d-l}$, and $f(x)$ is said to be *self-reciprocal* if $\tilde{f}(x) = ef(x)$ for some $e \in \mathbb{F}_{2^m}^\times$. Then by Equation (1) in Section 2, we have

$$x^{2n} - 1 = \tilde{f}_1(x)^2 \tilde{f}_2(x)^2 \dots \tilde{f}_r(x)^2.$$

Since $f_1(x), f_2(x), \dots, f_r(x)$ are pairwise coprime monic irreducible polynomials in $\mathbb{F}_{2^m}[x]$, $\tilde{f}_1(x), \tilde{f}_2(x), \dots, \tilde{f}_r(x)$ are pairwise coprime irreducible polynomials in $\mathbb{F}_{2^m}[x]$ as well. Hence for each integer j , $1 \leq j \leq r$, there is a unique integer j' , $1 \leq j' \leq r$, such that $\tilde{f}_j(x) = e_j f_{j'}(x)$ where $e_j \in \mathbb{F}_{2^m}^\times$. We assume that $f_j(x) = \sum_{l=0}^{d_j} c_l x^l$ where $c_j \in \mathbb{F}_{2^m}$. Then $x^{2d_j} f_j(x^{-1})^2 = \tilde{f}_j(x)^2$. From this, by Equation (4) and $x^{2n} = 1$ in $\mathcal{A}$ we deduce

$$\begin{aligned} \varrho(\varepsilon_j(x)) &= 1 - x^{2n - (\deg(b_j(x)) + 2d_j)} (x^{\deg(b_j(x))} v_j(x^{-1})) (x^{2d_j} f_j(x^{-1})^2) \\ &= 1 - x^{2n - (\deg(b_j(x)) + 2d_j)} \tilde{b}_j(x) \tilde{f}_j(x)^2 \\ &= 1 - h_j(x) f_{j'}(x)^2 \end{aligned}$$

where $h_j(x) = e_j^2 x^{2n - (\deg(b_j(x)) + 2d_j)} \tilde{b}_j(x) \in \mathcal{A}$. Similarly, by (4) it follows that $\varrho(\varepsilon_j(x)) = g_j(x) F_{j'}(x)$ for some $g_j(x) \in \mathcal{A}$. Then from these and by Equations (3) and (4) we deduce that $\varrho(\varepsilon_j(x)) = \varepsilon_{j'}(x)$.

As stated above, we see that for each integer j , $1 \leq j \leq r$, there is a unique integer j' , $1 \leq j' \leq r$, such that $\varrho(\varepsilon_j(x)) = \varepsilon_{j'}(x)$. We still use ϱ to denote this map $j \mapsto j'$; i.e., $\varrho(\varepsilon_j(x)) = \varepsilon_{\varrho(j)}(x)$. Whether ϱ denotes the ring automorphism of $\mathcal{A}$ or this map on the set $\{1, \dots, r\}$ is determined by context. The next lemma shows the compatibility of the two uses of ϱ .

Lemma 3.2 *With the notations above, we have the following:*

- (i) ϱ is a permutation on $\{1, \dots, r\}$ satisfying $\varrho^{-1} = \varrho$.
- (ii) After a rearrangement of $\varepsilon_1(x), \dots, \varepsilon_r(x)$ there are integers λ, ϵ such that $\varrho(j) = j$ for all $j = 1, \dots, \lambda$ and $\varrho(\lambda + l) = \lambda + \epsilon + l$ for all $l = 1, \dots, \epsilon$, where $\lambda \geq 1, \epsilon \geq 0$ and $\lambda + 2\epsilon = r$.
- (iii) For each integer j , $1 \leq j \leq r$, there is a unique invertible element e_j of $\mathbb{F}_{2^m}$ such that $\tilde{f}_j(x) = e_j f_{\varrho(j)}(x)$.
- (iv) For any integer j , $1 \leq j \leq r$, $\varrho(\varepsilon_j(x)) = \varepsilon_{\varrho(j)}(x)$ in the ring $\mathcal{A}$, and $\varrho(\mathcal{A}_j) = \mathcal{A}_{\varrho(j)}$. Then ϱ induces a ring isomorphism from $\mathcal{A}_j[u]/\langle u^k \rangle$ onto $\mathcal{A}_{\varrho(j)}[u]/\langle u^k \rangle$ in the natural way.

Proof. (i)–(iii) follow from the definition of the map ϱ , and (iv) follows from that $\mathcal{A}_j = \varepsilon_j(x)\mathcal{A}$ immediately. $\square$

Lemma 3.3 *Let $\alpha(x) = \sum_{j=1}^r \alpha_j(x)$, $\beta(x) = \sum_{j=1}^r \beta_j(x) \in \mathcal{A}[u]/\langle u^k \rangle$, where $\alpha_j(x), \beta_j(x) \in \mathcal{A}_j[u]/\langle u^k \rangle$. Then $\alpha(x)\varrho(\beta(x)) = \sum_{j=1}^r \alpha_j(x)\varrho(\beta_{\varrho(j)}(x))$.*

Proof. By Lemma 3.2 we have $\varrho(\beta_{\varrho(j)}(x)) \in \varrho(\mathcal{A}_{\varrho(j)}[u]/\langle u^k \rangle) = \mathcal{A}_{\varrho^2(j)}[u]/\langle u^k \rangle = \mathcal{A}_j[u]/\langle u^k \rangle$. Hence $\alpha_j(x)\varrho(\beta_{\varrho(j)}(x)) \in \mathcal{A}_j[u]/\langle u^k \rangle$ for all j . If $l \neq \varrho(j)$, then $j \neq \varrho(l)$ and $\mathcal{A}_j\mathcal{A}_{\varrho(l)} = \{0\}$ by Lemma 2.1(ii), which implies $\alpha_j(x)\varrho(\beta_l(x)) \in (\mathcal{A}_j[u]/\langle u^k \rangle)(\mathcal{A}_{\varrho(l)}[u]/\langle u^k \rangle) = \{0\}$. Hence $\alpha(x)\varrho(\beta(x)) = \sum_{j=1}^r \sum_{l=1}^r \alpha_j(x)\varrho(\beta_l(x)) = \sum_{j=1}^r \alpha_j(x)\varrho(\beta_{\varrho(j)}(x))$. $\square$

Lemma 3.4 *Using the notations above, for any $1 \leq j \leq r$ we have*

- (i) For any $\alpha = a(x) \in \mathcal{K}_j$, we denote $\hat{\alpha} = (\varphi_{\varrho(j)}^{-1}\varrho\varphi_j)(\alpha)$. Then $\hat{\cdot}$ is a ring isomorphism from $\mathcal{K}_j$ onto $\mathcal{K}_{\varrho(j)}$ such that the following diagram commutes

$$\begin{array}{ccc} \mathcal{K}_j = \mathbb{F}_{2^m}[x]/\langle f_j(x)^2 \rangle & \xrightarrow{\hat{\cdot}} & \mathcal{K}_{\varrho(j)} = \mathbb{F}_{2^m}[x]/\langle f_{\varrho(j)}(x)^2 \rangle \\ \varphi_j \downarrow & & \downarrow \varphi_{\varrho(j)} \\ \mathcal{A}_j & \xrightarrow{\varrho} & \mathcal{A}_{\varrho(j)} \end{array}$$

Specifically, we have $\hat{\alpha} = a(x^{-1}) \in \mathcal{K}_{\varrho(j)}$. Hence $\hat{f}_j(x) = f_j(x^{-1}) \in \mathcal{K}_{\varrho(j)}$.

- (ii) For any $\xi = \sum_{l=0}^{k-1} \alpha_l u^l \in \mathcal{K}_j[u]/\langle u^k \rangle$ where $\alpha_0, \alpha_1, \dots, \alpha_{k-1} \in \mathcal{K}_j$, define $\hat{\xi} = \sum_{l=0}^{k-1} \hat{\alpha}_l u^l$. Then ϱ induces a ring isomorphism from $\mathcal{K}_j[u]/\langle u^k \rangle$ onto $\mathcal{K}_{\varrho(j)}[u]/\langle u^k \rangle$ by the rule that $\varrho : \xi \mapsto \hat{\xi}$ ($\forall \xi \in \mathcal{K}_j[u]/\langle u^k \rangle$).

Proof (i) It follows from Lemma 2.1(iii) and Lemma 3.2 (iv).

(ii) It follows from (i) immediately. $\square$

Now, we give the dual code of each cyclic code over R of length $2n$.

Theorem 3.5 *Let $\mathcal{C}$ be a cyclic code over R of length $2n$ with canonical form decomposition $\mathcal{C} = \oplus_{j=1}^r \varepsilon_j(x)C_j$, where C_j is an ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$ given by Theorem 2.6. Then the dual code $\mathcal{C}^{\perp_E}$ of $\mathcal{C}$ is given by*

$$\mathcal{C}^{\perp_E} = \oplus_{j=1}^r \varepsilon_j(x)D_j,$$

where D_j is an ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$ given by one of the following eight cases for all $j = 1, \dots, r$:

case	C_j	$D_{\varrho(j)}$
1.	$\langle u^i \rangle$ ($0 \leq i \leq k$)	$\langle u^{k-i} \rangle$
2.	$\langle u^s f_j(x) \rangle$ ($0 \leq s \leq k-1$)	$\langle u^{k-s}, f_{\varrho(j)}(x) \rangle$
3.	$\langle u^i + u^t f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{i-t} \rangle)^\times$, $t \geq 2i-k$, $0 \leq t < i \leq k-1$)	$\langle u^{k-i} + u^{k+t-2i} f_{\varrho(j)}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{\varrho(j)}(x)}$
4.	$\langle u^i + f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $2i > k$ $0 < i \leq k-1$)	$\langle u^i + f_{\varrho(j)}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{\varrho(j)}(x)}$
5.	$\langle u^i + u^t f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $t < 2i-k$, $1 \leq t < i \leq k-1$)	$\langle u^{i-t} + f_{\varrho(j)}(x)\omega', u^{k-i} f_{\varrho(j)}(x) \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{\varrho(j)}(x)}$
6.	$\langle u^i, u^s f_j(x) \rangle$ ($0 \leq s < i \leq k-1$)	$\langle u^{k-s}, u^{k-i} f_{\varrho(j)}(x) \rangle$
7.	$\langle u^i + f_j(x)\omega, u^s f_j(x) \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^s \rangle)^\times$, $i+s \leq k-1$, $1 \leq s < i \leq k-1$)	$\langle u^{k-s} + u^{k-i-s} f_{\varrho(j)}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{\varrho(j)}(x)}$
8.	$\langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{s-t} \rangle)^\times$, $i+s \leq k+t-1$, $1 \leq t < s < i \leq k-1$)	$\langle u^{k-s} + u^{k+t-i-s} f_{\varrho(j)}(x)\omega', u^{k-i} f_{\varrho(j)}(x) \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{\varrho(j)}(x)}$

Proof Let $1 \leq j \leq r$, and E_j be an ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$ given by one of the following eight cases:

- (i) $E_j = \langle u^{k-i} \rangle$, if $C_j = \langle u^i \rangle$, where $0 \leq i \leq k$.
- (ii) $E_j = \langle u^{k-s}, f_j(x) \rangle$, if $C_j = \langle u^s f_j(x) \rangle$, where $0 \leq s \leq k-1$.
- (iii) $E_j = \langle u^{k-i} - u^{k+t-2i} f_j(x)\omega \rangle$, if $C_j = \langle u^i + u^t f_j(x)\omega \rangle$ where $\omega \in (\mathcal{F}_j[u]/\langle u^{i-t} \rangle)^\times$, $0 \leq t < i \leq k-1$ and $t \geq 2i-k$.
- (iv) $E_j = \langle u^i - f_j(x)\omega \rangle$, if $C_j = \langle u^i + f_j(x)\omega \rangle$ where $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $0 < i \leq k-1$ and $2i > k$.
- (v) $E_j = \langle u^{i-t} - f_j(x)\omega, u^{k-i} f_j(x) \rangle$, if $C_j = \langle u^i + u^t f_j(x)\omega \rangle$ where $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $1 \leq t < i \leq k-1$ and $t < 2i-k$.
- (vi) $E_j = \langle u^{k-s}, u^{k-i} f_j(x) \rangle$, if $C_j = \langle u^i, u^s f_j(x) \rangle$, where $0 \leq s < i \leq k-1$.
- (vii) $E_j = \langle u^{k-s} - u^{k-i-s} f_j(x)\omega \rangle$, if $C_j = \langle u^i + f_j(x)\omega, u^s f_j(x) \rangle$, where $\omega \in (\mathcal{F}_j[u]/\langle u^s \rangle)^\times$, $1 \leq s < i \leq k-1$, $i+s \leq k-1$.
- (viii) $E_j = \langle u^{k-s} - u^{k+t-i-s} f_j(x)\omega, u^{k-i} f_j(x) \rangle$, when $C_j = \langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$, where $\omega \in (\mathcal{F}_j[u]/\langle u^{s-t} \rangle)^\times$, $1 \leq t < s < i \leq k-1$, $i+s \leq k+t-1$.

Then by a direct calculation we deduce that $C_j E_j = \{a(x)b(x) \mid a(x) \in C_j, b(x) \in E_j\} = \{0\}$. Further, by Theorem 2.6 we have $|C_j||E_j| = 2^{2md_j k}$.

Since ϱ induces a ring isomorphism from $\mathcal{K}_j[u]/\langle u^k \rangle$ onto $\mathcal{K}_{\varrho(j)}[u]/\langle u^k \rangle$ by Lemma 3.4(ii), $\varrho(E_j)$ is an ideal of $\mathcal{K}_{\varrho(j)}[u]/\langle u^k \rangle$ and $|\varrho(E_j)| = |E_j|$. We denote $D_{\varrho(j)} = \varrho(E_j)$. Let $l = \varrho(j)$. Then $j = \varrho(l)$ by $\varrho^2 = \text{id}$. Now, set $\mathcal{D} = \sum_{j=1}^r \varepsilon_{\varrho(j)}(x) D_{\varrho(j)} = \oplus_{l=1}^r \varepsilon_l(x) D_l$, where $D_l = \varrho(E_{\varrho(l)})$. By Theorem 2.3, we see that $\mathcal{D}$ is a cyclic code over R of length $2n$.

For any $1 \leq j \leq r$, by Lemma 2.1(iii) we see that $\varepsilon_{\varrho(j)}(x) D_{\varrho(j)}$ is an ideal of $\mathcal{A}_{\varrho(j)}[u]/\langle u^k \rangle$. Moreover, we have $\varrho(D_{\varrho(j)}) = \varrho^2(E_j) = E_j$, and $\varrho(\varepsilon_{\varrho(j)}(x)) = \varepsilon_j(x)$ by Lemma 3.2(iv). From these and by Lemma 3.3, we deduce that

$$\begin{aligned} \mathcal{C} \cdot \varrho(\mathcal{D}) &= \left(\sum_{j=1}^r \varepsilon_j(x) C_j \right) \varrho \left(\sum_{l=1}^r \varepsilon_l(x) D_l \right) = \sum_{j=1}^r (\varepsilon_j(x) C_j) \varrho(\varepsilon_{\varrho(j)}(x) D_{\varrho(j)}) \\ &= \sum_{j=1}^r (\varepsilon_j(x) C_j) (\varepsilon_j(x) E_j) = \sum_{j=1}^r \varepsilon_j(x) (C_j E_j) = \{0\}, \end{aligned}$$

which implies $\mathcal{D} \subseteq \mathcal{C}^{\perp_E}$ by Lemma 3.1. On the other hand, by Lemma 2.1(iii) and Theorem 2.3(iii), we have $|\mathcal{C}| |\mathcal{D}| = (\prod_{j=1}^r |C_j|) (\prod_{j=1}^r |E_j|) = \prod_{j=1}^r (|C_j| |E_j|) = \prod_{j=1}^r 2^{md_j k} = 2^{2mk \sum_{j=1}^r d_j} = (2^{mk})^{2n} = |R[x]/\langle x^{2n} - 1 \rangle|$. Since $R = \mathbb{F}_{2^m}[u]/\langle u^k \rangle$ is a finite Frobenius ring, from the theory of linear codes over Frobenius rings (See [6], for example) we deduce that $\mathcal{C}^{\perp_E} = \mathcal{D}$.

Finally, we give the generator set of $D_{\varrho(j)} = \varrho(E_j)$, $1 \leq j \leq r$. By Lemma 3.2(iii), Lemma 3.4 and $x^{2n} = 1$, we have $\varrho(u^l) = u^l$ for all $1 \leq l \leq k$, $\varrho(\omega) = \widehat{\omega}$ for any $\omega \in \mathcal{K}_j[u]/\langle u^k \rangle$ and $\varrho(f_j(x)) = f_j(x^{-1}) = x^{2n-d_j} (x^{d_j} f_j(x^{-1})) = x^{2n-d_j} \widetilde{f}_j(x) = e_j x^{2n-d_j} f_{\varrho(j)}(x)$. Hence $\varrho(u^l f_j(x) \omega) = \varrho(u^l) \varrho(f_j(x)) \varrho(\omega) = u^l f_{\varrho(j)}(x) \omega'$, where $\omega' = e_j x^{2n-d_j} \widehat{\omega} \in \mathcal{F}_{\varrho(j)}[u]/\langle u^k \rangle$. Then the conclusions follows from (i)–(viii) immediately. $\square$

Finally, using Theorems 2.3, 2.6 and 3.5 we list all distinct self-dual cyclic codes over the ring R of length $2n$ by the following Theorem.

Theorem 3.6 *Using the notations in Lemma 3.2(ii), all distinct self-dual cyclic codes over the ring R of length $2n$ are give by:*

$$\mathcal{C} = \left(\oplus_{j=1}^{\lambda} \varepsilon_j(x) C_j \right) \oplus \left(\oplus_{j=\lambda+1}^{\lambda+\epsilon} (\varepsilon_j(x) C_j \oplus \varepsilon_{j+\epsilon}(x) C_{j+\epsilon}) \right),$$

where C_j is an ideal of $\mathcal{K}_j[u]/\langle u^k \rangle$ determined by the following conditions:

(A) If $1 \leq j \leq \lambda$, C_j is determined by the following conditions:

(A-i) When k is even, C_j is given by one of the following six cases:

(A-i-1) $C_j = \langle u^{\frac{k}{2}} \rangle$.

(A-i-2) $C_j = \langle f_j(x) \rangle$.

(A-i-3) $C_j = \langle u^{\frac{k}{2}+t} f_j(x) \omega \rangle$, where $0 \leq t \leq \frac{k}{2}-1$ and $\omega \in (\mathcal{F}_j[u]/\langle u^{\frac{k}{2}-t} \rangle)^{\times}$ satisfying $\omega + e_j x^{2n-d_j} \widehat{\omega} \equiv 0 \pmod{f_j(x)}$.

(A-i-4) $C_j = \langle u^i + f_j(x)\omega \rangle$, where $\frac{k}{2} + 1 \leq i \leq k-1$ and $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$ satisfying $\omega + e_j x^{2n-d_j} \widehat{\omega} \equiv 0 \pmod{f_j(x)}$.

(A-i-5) $C_j = \langle u^i, u^{k-i} f_j(x) \rangle$, where $\frac{k}{2} + 1 < i \leq k-1$.

(A-i-6) $C_j = \langle u^i + u^t f_j(x)\omega, u^{k-i} f_j(x) \rangle$, where $1 \leq t < k-i$, $\frac{k}{2} + 1 \leq i \leq k-1$ and $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i-t} \rangle)^\times$ satisfying $\omega + e_j x^{2n-d_j} \widehat{\omega} \equiv 0 \pmod{f_j(x)}$.

(A-ii) When k is odd, C_j is given by one of the following four cases:

(A-ii-1) $C_j = \langle f_j(x) \rangle$.

(A-ii-2) $C_j = \langle u^i + f_j(x)\omega \rangle$, where $\frac{k+1}{2} \leq i \leq k-1$ and $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$ satisfying $\omega + e_j x^{2n-d_j} \widehat{\omega} \equiv 0 \pmod{f_j(x)}$.

(A-ii-3) $C_j = \langle u^i, u^{k-i} f_j(x) \rangle$, where $\frac{k+1}{2} < i \leq k-1$.

(A-ii-4) $C_j = \langle u^i + u^t f_j(x)\omega, u^{k-i} f_j(x) \rangle$, where $1 \leq t < k-i$, $\frac{k+1}{2} \leq i \leq k-1$ and $\omega \in (\mathcal{F}_j[u]/\langle u^{k-i-t} \rangle)^\times$ satisfying $\omega + e_j x^{2n-d_j} \widehat{\omega} \equiv 0 \pmod{f_j(x)}$.

(B) If $j = \lambda + l$ where $1 \leq l \leq \epsilon$, then $(C_j, C_{j+\epsilon})$ is given by one of the following eight cases:

case	C_j	$C_{j+\epsilon}$
B-1.	$\langle u^i \rangle$ ($0 \leq i \leq k$)	$\langle u^{k-i} \rangle$
B-2.	$\langle u^s f_j(x) \rangle$ ($0 \leq s \leq k-1$)	$\langle u^{k-s}, f_{j+\epsilon}(x) \rangle$
B-3.	$\langle u^i + u^t f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{i-t} \rangle)^\times$, $t \geq 2i-k, 0 \leq t < i \leq k-1$)	$\langle u^{k-i} + u^{k+t-2i} f_{j+\epsilon}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{j+\epsilon}(x)}$
B-4.	$\langle u^i + f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $2i > k, 0 < i \leq k-1$)	$\langle u^i + f_{j+\epsilon}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{j+\epsilon}(x)}$
B-5.	$\langle u^i + u^t f_j(x)\omega \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{k-i} \rangle)^\times$, $t < 2i-k, 1 \leq t < i \leq k-1$)	$\langle u^{i-t} + f_{j+\epsilon}(x)\omega', u^{k-i} f_{j+\epsilon}(x) \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{j+\epsilon}(x)}$
B-6.	$\langle u^i, u^s f_j(x) \rangle$ ($0 \leq s < i \leq k-1$)	$\langle u^{k-s}, u^{k-i} f_{j+\epsilon}(x) \rangle$
B-7.	$\langle u^i + f_j(x)\omega, u^s f_j(x) \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^s \rangle)^\times$, $i+s \leq k-1, 1 \leq s < i \leq k-1$)	$\langle u^{k-s} + u^{k-i-s} f_{j+\epsilon}(x)\omega' \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{j+\epsilon}(x)}$
B-8.	$\langle u^i + u^t f_j(x)\omega, u^s f_j(x) \rangle$ ($\omega \in (\mathcal{F}_j[u]/\langle u^{s-t} \rangle)^\times$, $i+s \leq k+t-1$, $1 \leq t < s < i \leq k-1$)	$\langle u^{k-s} + u^{k+t-i-s} f_{j+\epsilon}(x)\omega', u^{k-i} f_{j+\epsilon}(x) \rangle$ $\omega' = e_j x^{2n-d_j} \widehat{\omega} \pmod{f_{j+\epsilon}(x)}$

4 An Example

We consider cyclic codes of length 14 over $R = \mathbb{F}_2[u]/\langle u^4 \rangle = \mathbb{F}_2 + u\mathbb{F}_2 + u^2\mathbb{F}_2 + u^3\mathbb{F}_2$ ($u^4 = 0$) which is a finite chain ring of 16 elements.

In this case, we have $m = 1$, $k = 4$ and $n = 7$. It is known that $x^7 - 1 = f_1(x)f_2(x)f_3(x)$ where $f_1(x) = x+1$, $f_2(x) = x^3+x+1$ and $f_3(x) = x^3+x^2+1$ are irreducible polynomials in $\mathbb{F}_2[x]$ satisfying $\tilde{f}_1(x) = f_1(x)$ and $\tilde{f}_2(x) = f_3(x)$. Hence $r = 3$, $d_1 = 1$, $d_2 = d_3 = 3$, $\lambda = 1$ and $\epsilon = 1$.

By Corollary 2.7, the number of cyclic codes over R of length 14 is

$$\prod_{j=1}^3 N_{(2,d_j,4)} = \prod_{j=1}^3 (1 + 14 + \Omega_1(2^{d_j}, 4) + \Omega_2(2^{d_j}, 4) + (2^{d_j} - 1)\Gamma(2^{d_j}, 4)),$$

where $\Gamma(2^{d_j}, 4) = 1$ for all $j = 1, 2, 3$, $\Omega_1(2^{d_1}, 4) = 5$, $\Omega_2(2^{d_1}, 4) = 2$, $\Omega_1(2^{d_2}, 4) = 77$ and $\Omega_1(2^{d_3}, 4) = 14$ for $j = 2, 3$. Then we have $\prod_{j=1}^3 N_{(2,d_j,4)} = 23 \cdot 113^2 = 293687$. Precisely, by Theorem 2.6 all distinct cyclic codes over R of length 14 are given by

$$\mathcal{C} = \varepsilon_1(x)C_1 \oplus \varepsilon_2(x)C_2 \oplus \varepsilon_3(x)C_3 \text{ with } |\mathcal{C}| = |C_1||C_2||C_3|,$$

where

$\diamond \varepsilon_1(x) = x^{12} + x^{10} + x^8 + x^6 + x^4 + x^2 + 1$, $\varepsilon_2(x) = x^8 + x^4 + x^2 + 1$ and $\varepsilon_3(x) = x^{12} + x^{10} + x^6 + 1$.

$\diamond C_1$ is an ideal of the ring $\mathcal{K}_1[u]/\langle u^4 \rangle$, where $\mathcal{K}_1 = \mathbb{F}_2[x]/\langle (x+1)^2 \rangle$ and $\mathcal{F}_1 = \mathbb{F}_2[x]/\langle x+1 \rangle = \mathbb{F}_2$, given by the following table:

case	number	C_1	$ C_1 $
I.	5	$\langle u^i \rangle \ (0 \leq i \leq 4)$	4^{4-i}
II.	4	$\langle u^s(x+1) \rangle \ (0 \leq s \leq 3)$	2^{4-s}
III.	5	$\langle u + (x+1) \rangle$	4^3
		$\langle u^2 + (x+1) \rangle, \langle u^2 + (x+1)(1+u) \rangle, \langle u^2 + u(x+1) \rangle$	4^2
		$\langle u^3 + u^2(x+1) \rangle$	4
IV.	2	$\langle u^3 + (x+1) \rangle$	2^4
		$\langle u^3 + u(x+1) \rangle$	2^3
V.	6	$\langle u^i, u^s(x+1) \rangle \ (0 \leq s < i \leq 3)$	$2^{8-(i+s)}$
VI.	1	$\langle u^2 + (x+1), u(x+1) \rangle$	2^5

$\diamond$ For $j = 2, 3$, C_j is an ideal of $\mathcal{K}_j[u]/\langle u^4 \rangle$, where $\mathcal{K}_j = \mathbb{F}_2[x]/\langle f_j(x)^2 \rangle$ and $\mathcal{F}_j = \mathbb{F}_2[x]/\langle f_j(x) \rangle = \{a_0 + a_1x + a_2x^2 \mid a_0, a_1, a_2 \in \mathbb{F}_2\}$, given by the following table:

case	number	C_j	$ C_j $
I.	5	$\bullet \langle u^i \rangle \ (0 \leq i \leq 4)$	64^{4-i}
II.	4	$\bullet \langle u^s f_j(x) \rangle \ (0 \leq s \leq 3)$	8^{4-s}
III.	7	$\bullet \langle u + f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	64^3
	56	$\bullet \langle u^2 + f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2 + u(b_0 + b_1x + b_2x^2), (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2, b_0, b_1, b_2 \in \mathbb{F}_2)$	64^2
	7	$\bullet \langle u^2 + u f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	64^2
	7	$\bullet \langle u^3 + u^2 f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	64
IV.	7	$\bullet \langle u^3 + f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	8^4
	7	$\bullet \langle u^3 + u f_j(x)\omega \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	8^3
V.	6	$\bullet \langle u^i, u^s f_j(x) \rangle \ (0 \leq s < i \leq 3)$	$8^{8-(i+s)}$
VI.	7	$\bullet \langle u^2 + f_j(x)\omega, u f_j(x) \rangle \ (\omega = a_0 + a_1x + a_2x^2, (a_0, a_1, a_2) \neq (0, 0, 0), a_0, a_1, a_2 \in \mathbb{F}_2)$	8^5

Moreover, by Theorem 3.6 all distinct self-dual cyclic codes over R of length 14 are given by $\mathcal{C} = \varepsilon_1(x)C_1 \oplus \varepsilon_2(x)C_2 \oplus \varepsilon_3(x)C_3$, where

- $\bullet C_1$ is given by one of the following cases:
- (A-i-1) $C_1 = \langle u^2 \rangle$.

- (A-i-2) $C_1 = \langle (x+1) \rangle$.
 (A-i-3) $C_1 = \langle u^2 + (x+1) \rangle$, $C_1 = \langle u^2 + (x+1)(1+u) \rangle$; $C_1 = \langle u^2 + u(x+1) \rangle$.
 (A-i-4) $C_1 = \langle u^3 + (x+1) \rangle$.
 (A-i-5) $C_1 = \langle u^3, u(x+1) \rangle$.
 • (C_2, C_3) is given by one pair of ideals in the following table:

case	C_2	C_3
B-1.	$\langle u^i \rangle, 0 \leq i \leq 4$	$\langle u^{k-i} \rangle$
B-2.	$\langle u^s f_2(x) \rangle, 0 \leq s \leq 3$	$\langle u^{4-s}, f_3(x) \rangle$
B-3.	$\langle u + f_2(x)\omega_1 \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u^3 + u^2 f_3(x)\omega'_1 \rangle, \omega'_1 = x^{11}\hat{\omega}_1$
	$\langle u^2 + f_2(x)\omega_2 \rangle, \omega_2 \in (\mathcal{F}_2[u]/\langle u^2 \rangle)^\times$	$\langle u^2 + f_3(x)\omega'_2 \rangle, \omega'_2 = x^{11}\hat{\omega}_2$
	$\langle u^2 + u f_2(x)\omega_1 \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u^2 + u f_3(x)\omega'_1 \rangle, \omega'_1 = x^{11}\hat{\omega}_1$
	$\langle u^3 + u^2 f_2(x)\omega_1 \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u + f_3(x)\omega'_1 \rangle, \omega'_1 = x^{11}\hat{\omega}_1$
B-4.	$\langle u^3 + f_2(x)\omega_1 \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u^3 + f_3(x)\omega'_1 \rangle, \omega'_1 = x^{11}\hat{\omega}_1$
B-5.	$\langle u^3 + u f_2(x)\omega_1 \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u^2 + f_3(x)\omega'_1, u f_3(x) \rangle, \omega'_1 = x^{11}\hat{\omega}_1$
B-6.	$\langle u^i, u^s f_2(x) \rangle, 0 \leq s < i \leq 3$	$\langle u^{4-s}, u^{4-i} f_3(x) \rangle$
B-7.	$\langle u^2 + f_2(x)\omega_1, u f_2(x) \rangle, \omega_1 \in \mathcal{F}_2^\times$	$\langle u^3 + u f_3(x)\omega'_1 \rangle, \omega'_1 = x^{11}\hat{\omega}_1$

in which $\omega_1 = a_0 + a_1x + a_2x^2$, $(a_0, a_1, a_2) \neq (0, 0, 0)$, $a_0, a_1, a_2 \in \mathbb{F}_2$, $\omega_2 = a_0 + a_1x + a_2x^2 + u(b_0 + b_1x + b_2x^2)$, $(a_0, a_1, a_2) \neq (0, 0, 0)$, $a_0, a_1, a_2, b_0, b_1, b_2 \in \mathbb{F}_2$, $\omega'_1 = x^{11}(a_0 + a_1x^{-1} + a_2x^{-2}) \pmod{f_3(x)}$ and $\omega'_2 = x^{11}(a_0 + a_1x^{-1} + a_2x^{-2} + u(b_0 + b_1x^{-1} + b_2x^{-2})) \pmod{f_3(x)}$.

Therefore, the number of self-dual cyclic codes of length 14 over $\mathbb{F}_2 + u\mathbb{F}_2 + u^2\mathbb{F}_2 + u^3\mathbb{F}_2$ ($u^4 = 0$) is equal to $7 \cdot 113 = 791$.

5 Conclusions and further research

We have developed a theory for cyclic codes of length $2n$ over the finite chain ring $R = \mathbb{F}_{2^m}[u]/\langle u^k \rangle = \mathbb{F}_{2^m} + u\mathbb{F}_{2^m} + \dots + u^{k-1}\mathbb{F}_{2^m}$ ($u^k = 0$) for any integer $k \geq 2$ and positive odd integer n , including the enumeration and construction of these codes, the dual code and self-duality for each of these codes. These codes enjoy a rich algebraic structure compared to arbitrary linear codes (which makes the search process much simpler). Obtaining some bounds for minimal distance such as BCH-like of a cyclic code over the ring R by just looking at the representation of such codes are future topics of interest.

Acknowledgements Part of this work was done when Yonglin Cao was visiting Chern Institute of Mathematics, Nankai University, Tianjin, China. Yonglin Cao would like to thank the institution for the kind hospitality. This research is supported in part by the National Key Basic Research Program of China (Grant No. 2013CB834204) and the National Natural Science Foundation of China (Grant No. 11471255).

References

1. T. Abualrub, I. Siap, Cyclic codes over the ring $\mathbb{Z}_2 + u\mathbb{Z}_2$ and $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2$, Des. Codes Cryptogr. **42** (2007), 273–287.
2. M. Al-Ashker, M. Hamoudeh, Cyclic codes over $\mathbb{Z}_2 + u\mathbb{Z}_2 + u^2\mathbb{Z}_2 + \dots + u^{k-1}\mathbb{Z}_2$, Turk. J. Math. **35**(4) (2011), 737–749.

3. A. Bonnetcaze, P. Udaya, Cyclic codes and self-dual codes over $\mathbb{F}_2 + u\mathbb{F}_2$, IEEE Trans. Inform. Theory **45** (1999), 1250–1255.
4. H. Q. Dinh, Constacyclic codes of length 2^s over Galois extension rings of $\mathbb{F}_2 + u\mathbb{F}_2$, IEEE Trans. Inform. Theory **55** (2009), 1730–1740.
5. H. Q. Dinh, Constacyclic codes of length p^s over $\mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$, J. Algebra **324** (2010), 940–950.
6. S. T. Dougherty, J-L. Kim, H. Kulosman, H. Liu: Self-dual codes over commutative Frobenius rings, Finite Fields Appl. **16**, 14–26 (2010).
7. G. Norton, A. Sălăgean-Mandache, On the structure of linear and cyclic codes over finite chain rings, Appl. Algebra in Engrg. Comm. Comput. **10** (2000), 489–506.
8. A. K. Singh, P. K. Kewat, On cyclic codes over the ring $\mathbb{Z}_p[u]/\langle u^k \rangle$, Des. Codes Cryptogr. **72** (2015), 1–13.

